



École nationale
de la statistique
et de l'analyse
de l'information

EVALUATION OF THE EXCHANGEABILITY NON-EXCHANGEABILITY METHODOLOGY IN PHASE II POC STUDIES

Audrey PUAUD

20nd November 2018



Context

EXNEX methodology

Application: phase II POC

Discussion



What?

Oncology

Early phases (I/II)


Multiple **subgroups**



Why?

To **identify** interesting strata **earlier**

To **combine information** in a single trial





What?

Oncology

Early phases (I/II)


Multiple subgroups



Why?


To **identify** interesting strata **earlier**

To **combine information** in a single trial






- **Phase II Proof-of-Concept** trials
- **Binary** endpoint: response vs. no response (e.g tumour shrinkage)





Stratification modeling

$$\theta_j \sim N(m_j, v_j)$$



Estimates per group




Reliability?
Over-interpretation?



Pooling

$$\theta \sim N(m, v)$$



Precision




No estimate per group
Accuracy?

Hierarchical modeling

$$\theta_j | \mu, \tau \sim N(\mu, \tau^2)$$

$$\mu \sim N(m, v) \quad \tau \sim HN(\sigma^2)$$



Estimates per group
Borrowing information



Exchangeability assumption?



Neuenschwander, B., Wandel, S., Roychoudhury, S. & Bailey, S. (2016). Robust exchangeability designs for early phase clinical trials with multiple strata.

Robust mixture extension of the standard hierarchical approach

EX: Exchangeability across similar strata

+

NEX: Non-exchangeability for extreme strata



EXNEX

EXNEX approach


Mixture of hierarchical and stratified modeling method

EXCHANGEABLE (EX)

$$\theta_j | \mu, \tau \sim N(\mu, \tau^2)$$

$$\mu \sim N(m, v)$$


$$\tau \sim HN(\sigma^2)$$



With probability w_j

NON-EXCHANGEABLE (NEX)

$$\theta_j \sim N(m_j, v_j)$$



With probability $1 - w_j$



Bayesian inference:

$$\text{Prior} \times \text{Likelihood} = \boxed{\text{Posterior}} \quad \bigcirc ?$$


Bayesian inference:

$$\text{Prior} \times \text{Likelihood} = \text{Posterior}$$



Markov chain Monte Carlo (**MCMC**) algorithms

Rjags (Gibbs sampler)




$$\theta_j | \mu_j, \tau_j, i_j \sim N(\mu_j, \tau_j^2)$$

$$i_j \sim \text{Cat}(w_j)$$

For each MCMC iteration:

$$\begin{cases} \mu_j = \mu & \text{et} \quad \tau_j = \tau & \text{if } i_j = 1 \\ \mu_j = m_j & \text{et} \tau_j = \sqrt{v_j} & \text{if } i_j = 2 \end{cases}$$



$$\theta_j | \mu_j, \tau_j, i_j \sim N(\mu_j, \tau_j^2)$$

$$i_j \sim \text{Cat}(w_j)$$

For each MCMC iteration:

$$\left\{ \begin{array}{l} \mu_j = \mu^{(1)} \text{ et } \tau_j = \tau^{(1)} \text{ if } i_j = 1 \\ \mu_j = \mu^{(2)} \text{ et } \tau_j = \tau^{(2)} \text{ if } i_j = 2 \\ \mu_j = m_j \text{ et } \tau_j = \sqrt{v_j} \text{ if } i_j = 3 \end{array} \right.$$

For each MCMC iteration

Strata j

EXCHANGEABLE


- Follows one of the **common** EX distribution(s)
- **Shares information** with the other exchangeable subgroups of this iteration

NON-EXCHANGEABLE

- Follows its **own** NEX distribution
- **No sharing** of information with the other subgroups




- Phase II POC study (binary endpoint) with 4 subgroups:



- Success criteria for each strata:

$$\hat{\pi} \geq 20\% \ \& \ P(\pi > 10\%) \geq 81\% \text{ or } 91\%$$

- Comparison of **hierarchical**, **stratified** and **EXNEX** modeling methods



EXNEX model:

$$w_j = p_j = (0.25, 0.25, 0.50)$$

EX component:

$$\mu^{(1)} \sim N(m = \text{logit}(0.1), v = 10.1)$$


$$\tau^{(1)} \sim HN(\sigma^2 = 1^2)$$

$$\mu^{(2)} \sim N(m = \text{logit}(0.3), v = 3.8)$$

$$\tau^{(2)} \sim HN(\sigma^2 = 1^2)$$

NEX component:

$$\theta_j \sim N(m_j = \text{logit}(0.2), v_j = 6.3)$$



■ Hierarchical model:

$$\mu \sim N(m = \text{logit}(0.2), v = 5.3)$$

$$\tau \sim HN(\sigma^2 = 1^2)$$

■ Stratified model:

$$\theta_j \sim N(m_j = \text{logit}(0.2), v_j = 6.3)$$

- 2 scenarii:

Homogeneous



Nugget scenario






Context

EXNEX

Application

Discussion



Homogeneous situation

Priors:
 $\hat{m}^1 = \text{logit}(0.1)$
 $\hat{m}^2 = \text{logit}(0.3)$
 $\hat{m}_j = \text{logit}(0.2)$
 $\hat{w}_j = (0.25, 0.25, 0.50)$

Context

EXNEX

Application

Discussion

True
response rate

30%

35%

40%

30%

Posterior distributions				
Strata	1	2	3	4
EX component				
\hat{m}^1	logit(0.20)			
\hat{w}_j^1	0.27	0.28	0.27	0.27
\hat{m}^2	logit(0.32)			
\hat{w}_j^2	0.38	0.38	0.37	0.36
NEX component				
\hat{m}_j	logit(0.23)	logit(0.24)	logit(0.25)	logit(0.22)
\hat{w}_j^3	0.35	0.34	0.37	0.37



Priors:

$\hat{m}^1 = \text{logit}(0.1)$

$\hat{m}^2 = \text{logit}(0.3)$

$\hat{m}_j = \text{logit}(0.2)$

$\hat{w}_j = (0.25, 0.25, 0.50)$




Context

EXNEX

Application

Discussion





Context

EXNEX

Application

Discussion

True
response rate

65%

35%

40%

30%

Posterior distributions				
Strata	1	2	3	4
EX component				
\hat{m}^1	logit(0.24)			
\hat{w}_j^1	0.22	0.26	0.26	0.25
\hat{m}^2	logit(0.38)			
\hat{w}_j^2	0.31	0.34	0.35	0.33
NEX component				
\hat{m}_j	logit(0.39)	logit(0.25)	logit(0.26)	logit(0.23)
\hat{w}_j^3	0.47	0.40	0.38	0.42




Priors:

$$\hat{m}^1 = \text{logit}(0.1)$$

$$\hat{m}^2 = \text{logit}(0.3)$$

$$\hat{m}_j = \text{logit}(0.2)$$

$$\hat{w}_j = (0.25, 0.25, 0.50)$$





Context

EXNEX

Application

Discussion

- **42** scenarii
- **2 000** simulated clinical trials for each scenario
- **2, 3, 4 or 6** subgroups
- Several **degrees of heterogeneity** inter-strata
- **3 metrics**: probability of success, bias and MSE



Context

EXNEX

Application

Discussion




Context

EXNEX

Application

Discussion




Context

EXNEX

Application

Discussion





Context

EXNEX

Application

Discussion

BAYESIAN
MIXTURE
MODEL


GOOD
COMPROMISE

CHALLENGING
PRIOR
SPECIFICATIONS

PLUS-VALUE
WITH ≥ 3
SUBGROUPS



EXCHANGEABILITY
ASSUMPTION



Thank you for your attention!

Any questions?






École nationale
de la statistique
et de l'analyse
de l'information

EVALUATION OF THE EXCHANGEABILITY NON-EXCHANGEABILITY METHODOLOGY IN PHASE II POC STUDIES

Audrey PUAUD
20nd November 2018




■ Probability of success in stratified model (logit(0.2)):






- Success criteria for each strata:

$$\hat{\pi} \geq 20\% \quad \& \quad P(\pi > 10\%) \geq 80\% \text{ or } 90\%$$



Borderline cases in stratified analyses





- **Prior specifications:** weakly informative setting (1)

$$\theta_j = \log\left(\frac{\pi_j}{1 - \pi_j}\right)$$

$$\theta_j \sim N(m_j, v_j)$$

$$\mu \sim N(m, v)$$

$$\tau \sim HN(1^2)$$

$$v_j = V(\theta_j) = \frac{1}{p} + \frac{1}{1 - p}$$

One observation

$$v = \underbrace{V(\theta_j)}_{\text{Between variance}} - \underbrace{E(\tau^2)}_{\text{Within variance}} = V(\theta_j) + E(\tau)^2 - V(\tau)$$

*Between
variance*

*Within
variance*



- **Prior specifications:** weakly informative setting (2)

$$\theta_j = \log\left(\frac{\pi_j}{1 - \pi_j}\right)$$

$$\theta_j \sim N(m_j, v_j)$$

$$\mu \sim N(m, v)$$

$$\tau \sim HN(1^2)$$

$$v_j = V(\theta_j) = \frac{1}{0.2} + \frac{1}{1 - 0.2} = \mathbf{6.3}$$

One observation

$$\begin{aligned} v &= V(\theta_j) - E(\tau^2) = V(\theta_j) + E(\tau)^2 - V(\tau) \\ &= 6.3 + 0 - 1^2 = \mathbf{5.3} \end{aligned}$$



■ Gibbs sampling

Iterative algorithm:

1. Initialization of the p parameters: $\theta^0 = (\theta_1^0, \dots, \theta_p^0)$
2. Updating (t^{th} iteration):

$$\begin{aligned}\theta_1^t &= \pi_1(\cdot | \theta_2^{t-1}, \dots, \theta_p^{t-1}) \\ \theta_2^t &= \pi_2(\cdot | \theta_1^t, \theta_3^{t-1}, \dots, \theta_p^{t-1}) \\ &\vdots \\ \theta_p^t &= \pi_p(\cdot | \theta_1^t, \dots, \theta_{p-1}^t)\end{aligned}$$



■ Case studies: homogeneous scenario

True response rate 30% 35% 40% 30%

Posterior distributions (median)				
Strata	1	2	3	4
EX component				
\hat{m}^1	logit(0.28)			
\hat{m}^2	logit(0.32)			
NEX component				
\hat{m}_j	logit(0.26)	logit(0.29)	logit(0.31)	logit(0.25)



■ Case studies: heterogeneous scenario

True response rate **65%** **35%** **40%** **30%**

Posterior distributions (median)				
Strata	1	2	3	4
EX component				
\hat{m}^1	logit(0.33)			
\hat{m}^2	logit(0.39)			
NEX component				
\hat{m}_j	logit(0.56)	logit(0.30)	logit(0.32)	logit(0.25)