

# Scale Loss Score (SLoS): a novel measure of drug benefit-risk assessment

**Gaelle Saint-Hilary<sup>1,2</sup>,**

V. Robert<sup>1</sup>, M. Gasparini<sup>2</sup>, T. Jaki<sup>3</sup> and P. Mozgunov<sup>3</sup>

<sup>1</sup>Servier, France ; <sup>2</sup>Polytechnic University of Turin, Italy ;

<sup>3</sup>Lancaster University, UK


*gaelle.saint-hilary@servier.com*

SFdS B&S, 20 November 2018

- **MultiCriteria Decision Analysis (MCDA)** is a popular quantitative method to assess the **benefit-risk (BR) balance** of treatments: it permits to summarize the benefits and the risks of a drug in a **single utility score**
- The utility score is often derived using a **linear model** which might lead to **counter-intuitive conclusions**, for example, a recommendation of a non-effective drug
- We propose **Scale Loss Score (SLoS)** as a new tool for benefit-risk assessment: it is based on strong theoretical principles, addresses the issues of the linear MCDA model and can lead to more **meaningful recommendations**

- **Introduction**
- **Linear MultiCriteria Decision Analysis (MCDA)**
  - Model
  - Counter-intuitive conclusions
- **Scale Loss Score (SLoS)**
  - Model
  - Differences with Linear MCDA
  - Weight mapping
  - Case-study: telithromycin
  - Simulation study
- **Conclusion**

# Linear Multi-Criteria Decision Analysis (MCDA)



**MCDA linear utility score**

$$u(\xi_i, \mathbf{w}) := \sum_{j=1}^n w_j u_j(\xi_{ij})$$

**Higher utility score  $\rightarrow$  more preferable BR balance**


# Linear Multi-Criteria Decision Analysis (MCDA)

## Partial value functions


$u_j(\xi_{ij})$ : linear partial value functions - map the performances on criterion  $j$  to a  $(0, 1)$  scale

$$u_j(\xi_{ij}) = \frac{\xi_{ij} - \xi'_{ij}}{\xi''_{ij} - \xi'_{ij}}, \quad \xi'_{ij} \text{ and } \xi''_{ij} \text{ the worst and best values}$$

**The greater the criterion value the better**




**The lower the criterion value the better**



# Linear Multi-Criteria Decision Analysis (MCDA)

## Example with two criteria

Contours of  $1 - u(\xi_{i1}, \xi_{i2}, w)$




# Linear Multi-Criteria Decision Analysis (MCDA)

Might lead to counter-intuitive conclusions

(1) The benefit-risk trade-off is the same for all levels of risk/benefit

Contours of  $1 - u(\xi_{i1}, \xi_{i2}, w = 0.25)$




- Drug B increases the benefit from 0.15 to 0.30 compared to A ( $\times 2$ )
- An increase of +5% in risk could be tolerated to have the same utility

# Linear Multi-Criteria Decision Analysis (MCDA)

Might lead to counter-intuitive conclusions

(1) The benefit-risk trade-off is the same for all levels of risk/benefit

Contours of  $1 - u(\xi_{i1}, \xi_{i2}, w = 0.25)$



- Drug D increases the benefit from 0.80 to 0.95 compared to C ( $\times 1.1875$ )
- An increase of +5% in risk could be tolerated to have the same utility

The same benefit increase is not as relatively large, only a smaller increase in risk may usually be tolerated



# Linear Multi-Criteria Decision Analysis (MCDA)

Might lead to counter-intuitive conclusions

(2) Drugs with no benefit or extreme risk can be recommended

2 criteria, fixed parameter values and  $w = 0.25$

	Example 1		Example 2	
	Low benefit and risk		High benefit and risk	
	Drug 1	Drug 2	Drug 1	Drug 2
Benefit	0%	30%	96%	50%
Risk	9%	20%	100%	85%
MCDA utility	0.6825	0.6750	0.2400	0.2375

Even if none of those drugs are likely to be taken to the market, these examples reveal some counter-intuitive conclusions

# Linear Multi-Criteria Decision Analysis (MCDA)

Might lead to counter-intuitive conclusions


With Linear MCDA:

- (1) The benefit–risk trade-off is the same for all levels of risk/benefit
- (2) Drugs with no benefit or extreme risk can be recommended

We advocate two properties of a benefit–risk analysis measure:

- For a given increase in benefit, one can tolerate a larger increase in risk if the amount of benefit is small than if it is high  
→ Convex preferences (i.e. concavity of equal loss contours)
- One is not interested in the level of risk (benefit) if the drug does not treat (harm all) patients  
→ Strong penalisation of extreme low benefit values and extreme high risk values

# Scale Loss Score (SLoS)




$$l(\xi_i, \tilde{w}) := \sum_{j=1}^n \left( \frac{1}{u_j(\xi_{ij})} \right)^{\tilde{w}_j}$$

Lower loss score  $\rightarrow$  more preferable BR balance

## Example with two criteria


Contours of  $l(\xi_{i1}, \xi_{i2}, \tilde{w})$



# Scale Loss Score (SLoS)

Addresses the counter-intuitive conclusions of Linear MCDA

Contours of  $I(\xi_{i1}, \xi_{i2}, \tilde{w} = 0.25)$



- Drug B increases the benefit from 0.15 to 0.30 ( $\times 2$ ) compared to A
- Drug D increases the benefit from 0.80 to 0.95 ( $\times 1.1875$ ) compared to C

For a given increase in benefit, a **smaller** increase in risk is tolerated if benefit is high than if it is low

# Scale Loss Score (SLoS)

Addresses the counter-intuitive conclusions of Linear MCDA

2 criteria, fixed parameter values and  $w = \tilde{w} = 0.25$

	Example 1		Example 2	
	Low benefit and risk		High benefit and risk	
	Drug 1	Drug 2	Drug 1	Drug 2
Benefit	0%	30%	96%	50%
Risk	9%	20%	100%	85%
MCDA utility	0.6825	0.6750	0.2400	0.2375
SLoS	$+\infty$	2.5334	$+\infty$	5.3381

→ SLoS strongly penalizes extremely low benefit values and extremely high risk values

Drugs with no benefit or extreme risk **can never be** recommended

# Weight mapping

Comprehensive work was already published or is on-going on weight elicitation for MCDA


→ We propose a simple way to map MCDA weights  $w_j$  to SLoS weights  $\tilde{w}_j$

With two criteria:

$$\frac{\tilde{w}_j}{1 - \tilde{w}_j} \cdot 2^{2\tilde{w}_j - 1} = \frac{w_j}{1 - w_j},$$


No analytical solution, but  $\tilde{w}_j$  can be approximated by line search

## Weight mapping



# Case study: telithromycin

IMI PROTECT Benefit-Risk Group example



## Proba(telithromycin > $\beta$ -lactam antibiotics)

	Community Acquired Pneumonia (CAP)	Acute Bacterial Sinusitis (ABS)
MCDA	59%	71%
SLoS	51%	55%

→ SLoS results are more in line with the regulatory authorities concerns on ABS indication



# Simulations (1/2)


Comprehensive simulation study investigating the performances of MCDA and SLoS in many different scenarios

## Simulation scenarios with two criteria

Probability of Risk $\theta_{i2}$	Probability of Benefit $\theta_{i1}$								
	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.8	0.9
0.9	◊●	◊	◊	◊	◊	◊	◊	◊	◊●
0.8	◊	◊	◊	◊	◊	◊	◊	◊	◊
0.7	◊	◊	◊●	◊	◊	◊	◊●	◊	◊
0.6	◊	◊	◊	◊	◊	◊	◊	◊	◊
0.5	◊	◊	◊	◊	◊●	◊	◊	◊	◊
0.4	◊	◊	◊	◊	◊	◊	◊	◊	◊
0.3	◊	◊	◊	◊	◊	◊	◊	◊	◊
0.2	◊	◊	◊	◊	◊	◊	◊	◊	◊
0.1	◊●	◊	◊	◊	◊	◊	◊	◊	◊

● = treat

## Simulation results with two




Number of criteria, imbalanced weights, correlated criteria, impact of the weight mapping...

## Simulation scenarios with four criteria

Probability of Risk ( $\theta_{i3}$ and $\theta_{i4}$ )		Probability of Benefit ( $\theta_{i1}$ and $\theta_{i2}$ )								
		0.1			0.5			0.9		
		0.1	0.5	0.9	0.1	0.5	0.9	0.1	0.5	0.9
0.9	0.9	◊●	◊	◊	◊	◊	◊	◊	◊	◊●
	0.5	◊	◊●	◊	◊	◊	◊	◊●	◊	◊
	0.1	◊	◊	◊	◊	◊	◊	◊	◊	◊
0.5	0.9	◊	◊	◊	◊	◊	◊	◊	◊	◊
	0.5	◊	◊	◊	◊●	◊	◊	◊	◊	◊
	0.1	◊	◊	◊	◊	◊	◊	◊	◊	◊
0.1	0.9	◊	◊	◊	◊	◊	◊	◊	◊	◊
	0.5	◊●	◊	◊	◊	◊	◊	◊	◊●	◊
	0.1	◊	◊	◊	◊	◊	◊	◊	◊	◊

## Simulation results with four criteria



Comprehensive simulation study investigating the performances of MCDA and SLoS in many different scenarios

## **Main conclusions from the simulation study:**

- Both are robust to correlations between outcomes
- Similar conclusions in many cases
- Clear advantage of SLoS when drugs have no benefit or extreme risk

- **Scale Loss Score (SLoS)** offers the same advantages as Linear MCDA to summarise the drug benefit–risk balance in a **single measure**
  - SLoS has additional **desirable properties**:
    - Avoids recommendations of non-effective or extremely unsafe drugs
    - Tolerates larger increases in risk for a given increase in benefit when the amount of benefit is small than when it is high
- **Better reflects human's natural preferences**
- Alternative approach: handling non-constant trade-offs by varying the shape of the partial value functions (non-linear MCDA)
    - Non-trivial approach, and eliciting non-linear partial value functions is very challenging

# Main references I

Saint-Hilary G, Robert V, Gasparini M, Jaki T and Mozgunov P. A novel measure of drug benefit-risk assessment based on scale loss score. *Stat Methods Med Res*, 2018.

Saint-Hilary G, Cadour S, Robert V and Gasparini M. A simple way to unify multicriteria decision analysis (MCDA) and stochastic multicriteria acceptability analysis (SMAA) using a dirichlet distribution in benefit-risk assessment. *Biometrical Journal*, 59(3):567–578, 2017.

Mozgunov P, Jaki T and Gasparini M. Loss Functions in Restricted Parameter Spaces and Their Bayesian Applications. *arXiv:1706.02104*, 2017.

Mozgunov P and Jaki T. An information-theoretic approach for selecting arms in clinical trials. *arXiv:1708.02426*, 2017.

Berger JO. *Statistical decision theory and Bayesian Analysis (2nd ed.)*. Springer-Verlag, New York, 1985.

Mussen F, Salek S and Walker S. A quantitative approach to benefit-risk assessment of medicines - Part 1: the development of a new model using multi-criteria decision analysis. *Pharmacoepidemiology and Drug Safety*, 16:S2—S15, 2007.

Waddingham E, Mt-Isa S, Nixon R and Ashby D. A Bayesian approach to probabilistic sensitivity analysis in structured benefit-risk assessment. *Biometrical Journal*, 58:28–42, 2016.

# Main references II

Tervonen T, Gelhorn H, Sri Bhashyam S, Poon JL, Gries KS, Rentz A and Marsh K. MCDA swing weighting and discrete choice experiments for elicitation of patient benefit-risk preferences: a critical assessment. *Pharmacoepidemiology and Drug Safety*, pages 1–9, 2017.

Thokaka P, Devlin N, Marsh K, Baltussen R, Boysen M, Kalo Z, Longrenn T, Mussen F, Peacock S, Watkins J and IJzerman M. Multiple Criteria Decision Analysis for Health Care Decision Making - An Introduction: Report 1 of the ISPOR MCDA Emerging Good Practices Task Force. *Value in Health*, 19:1–13, 2016.

Marsh K, IJzerman M, Thokala P, Baltussen R, Boysen M, Kalo Z, Longrenn T, Mussen F, Peacock S, Watkins J and Devlin N. Multiple Criteria Decision Analysis for Health Care Decision Making - Emerging Good Practices: Report 2 of the ISPOR MCDA Emerging Good Practices Task Force. *Value in Health*, 19:125–137, 2016.

Langhans SD and Lienert J. Four common simplifications of multi-criteria decision analysis do not hold for river rehabilitation. *PLOS One*, 11(3):1–27, 03 2016.

Martin DM and Mazzotta M. Non-monetary valuation using Multi-Criteria Decision Analysis: Sensitivity of additive aggregation methods to scaling and compensation assumptions. *Ecosystem Services*, 29:13–22, 2018.

IMI PROTECT. Work package 5: Benefit-risk integration and representation. 2013. <http://www.imi-protect.eu/wp5.shtml>.

# Main references III

IMI PROTECT. Work package 5: Report 1:b:ii Benefit – Risk Wave 1 Case Study Report: Ketek<sup>®</sup> (telithromycin). 2012. <http://www.imi-protect.eu/documents/QuarteyetalBenefitRiskWave1CasestudyReportTelithromycinFeb2012.pdf>.

EMA. Benefit-risk methodology project. Work package 1 Report: description of the current practice of benefit-risk assessment for centralised procedure products in the EU regulatory network. 2011. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Report/2011/07/WC500109478.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Report/2011/07/WC500109478.pdf).

EMA. Benefit-risk methodology project. Work package 2 Report: Applicability of current tools and processes for regulatory benefit-risk assessment. 2010. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Report/2010/10/WC500097750.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Report/2010/10/WC500097750.pdf).

EMA. Benefit-risk methodology project. work package 3 report: Field tests. 2011. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Report/2011/09/WC500112088.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Report/2011/09/WC500112088.pdf).

EMA. Benefit-risk methodology project. Work package 4 Report: Benefit-risk tools and processes. 2012. [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Report/2012/03/WC500123819.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Report/2012/03/WC500123819.pdf).

**Thank-you!**